

# Fact Sheet for Assessing Decision-Making Capacity

## Relevance of Assessing Decision-Making Capacity (DMC)

Sometimes physicians, nurses, and others are called upon to determine if a patient has decision-making capacity and can consent to a treatment or to admission to a health care facility (e.g., a nursing home). This fact sheet helps them to make that determination.

The assessment of DMC in patient care is critically important. If the patient has not completed an advance directive, specified someone to make decisions for him or her, or discussed his/her preferences for medical treatment with others, then an assessment that the patient lacks DMC means that the patient will have lost control over the decisions that are made for him or her. In almost all circumstances, health caregivers are obligated to respect the wishes, i.e., informed consent or refusal, of a patient with DMC. If a patient has completed a proxy advance directive such as a durable power of attorney for health care (called health care proxy in NY or MA or medical power of attorney in WV), then an assessment that the patient lacks DMC means that the person designated in the patient's proxy advance directive assumes the authority to make decisions for the patient subject to any limitations the patient has specified in the authority or any specific instructions the patient has given for care. **See V below for questions to ask to assess if the patient has decision-making capacity.**

### I. Competency vs. Decision-making capacity (DMC)

Incompetency is determined by a court.

Decision-making capacity is a clinical judgment made by those authorized under state law to make the determination. In West Virginia, physicians, licensed psychologists, advanced nurse practitioners, and physician assistants can determine DMC.

### II. By law, patients are presumed to be competent.

### III. Mental Status Testing may be inadequate to determine DMC.

A patient may have deficits in one or more of the following and still have DMC:

- orientation
- attention span
- immediate recall
- long-term memory
- calculating ability

### IV. The capacity to make health care decisions requires that the patient have the following abilities:

- the ability to understand one's condition;

- the ability to appreciate the consequences (benefits and risks) of the main treatment options including non-treatment;
- the ability to judge the relationship between the treatment options and their consequences to one's values, preferences, and goals;
- the ability to reason and deliberate about one's options; and
- the ability to communicate one's decisions in a meaningful manner.

**V. In a practical sense, physicians, psychologists, nurses, social workers, and therapists can determine if a patient has decision-making capacity by whether the patient can give informed consent or refusal. The following questions are helpful to make this assessment:**

Can the patient understand what is wrong with him or her and what are the proposed procedures or treatments? Specifically, can the patient answer...

- As you understand it, what is your medical problem?
- How serious is your illness? What will happen if you are not treated?
- What do you think caused your illness, and why did it start when it did?

Can the patient understand the benefits and risks of the proposed procedure or treatment and the benefits and risks of the alternative procedures or treatments including non-treatment?

Is the patient able to reason and make a decision using the medical information which has been disclosed to him or her and to incorporate his or her personal values and wishes into the decision?

Is the patient able to explain why he or she made the health care decision that he or she did, and is the explanation consistent with his or her stated values and wishes?

**VI. Note that patients who are of advanced age, depressed, demented, or intellectually challenged may still possess DMC.**

**VII. DMC is not absolute or permanent**

Patients may possess sufficient understanding and reasoning ability to make some decisions (for example, agreeing to have blood drawn) but not others which require more complex levels of these skills (for example, the giving of advance directives). In this sense, DMC is said to be task-related and decision specific.

Patients may have DMC at one time but not another. If possible, decisions should be delayed until reversible problems that might alter DMC such as infection, electrolyte imbalance, or medication effect resolve.